

**CHLORINE DIOXIDE GAS FOR THE PREVENTION OF INFECTIOUS DISEASES**

Norio Ogata\*

R and D Center, Taiko Pharmaceutical Co., Ltd., Seikacho, Kyoto, Japan.

**\*Corresponding Author: Dr. Norio Ogata**

R and D Center, Taiko Pharmaceutical Co., Ltd., Seikacho, Kyoto, Japan.

Article Received on 20/06/2017

Article Revised on 10/07/2017

Article Accepted on 31/07/2017

**ABSTRACT**

The prevention of infectious diseases caused by infectious air-floating microbes in closed and semi-closed areas is strongly needed, but it is still difficult to achieve satisfactorily because of the absence of appropriate disinfection techniques. Fumigation is one of the methods to disinfect air contaminated with infectious microbes, but it absolutely requires evacuation of people from the areas to be disinfected because the gas concentration of disinfectant employed is usually very high and toxic to animals. Currently it is demonstrated that chlorine dioxide ( $\text{ClO}_2$ ) gas of extremely low concentrations that have no toxic effect to animals has strong anti-microbial activity against infectious microbes, such as bacteria and viruses. Disinfection using such low-concentration  $\text{ClO}_2$  gas does not require evacuation of people, and could be used to disinfect room air in the simultaneous presence of people. The use of  $\text{ClO}_2$  gas at very low concentrations may open new avenue of disinfection systems of room air without requiring evacuation of people. This review presents the details of the disinfection system of  $\text{ClO}_2$  gas.

**KEYWORDS:** Chlorine dioxide,  $\text{ClO}_2$ , Gas, Bacteria, Virus, Disinfection.**INTRODUCTION**

Chlorine dioxide ( $\text{ClO}_2$ ) is a liquid that exhibits dark orange color below  $11^\circ\text{C}$ .<sup>[1]</sup>  $\text{ClO}_2$  starts boiling above this temperature and produces yellow gas with a characteristic odor. It is a relatively stable free radical with one unpaired electron in its molecular orbital. Its molecular structure in a liquid state was demonstrated by an X-ray diffraction analysis.<sup>[2]</sup>  $\text{ClO}_2$  has long been used to disinfect tap water in place of chlorine in some countries. Contrary to chlorine disinfection,  $\text{ClO}_2$  disinfection does not produce potentially carcinogenic trihalomethane.<sup>[3]</sup> Microbes disinfected by  $\text{ClO}_2$  include bacteria, fungi, protozoa and viruses.<sup>[4]</sup>  $\text{ClO}_2$  dissolved in water and gaseous  $\text{ClO}_2$  are used to disinfect microbes. The detailed chemical dynamic mechanisms of inactivation of microbes by  $\text{ClO}_2$  are reported.<sup>[5]</sup> Currently, extremely low concentrations of  $\text{ClO}_2$  gas, on the order of 0.01- 0.05 ppm (parts per million) (volume ratio), are used to disinfect microbes in room air. The gas still possesses antimicrobial activity at these low levels.<sup>[6,7]</sup> Such extremely low levels of the gas are reported to be non-hazardous to animals,<sup>[8]</sup> and its potential use in closed and semi-closed spaces without a need of evacuation of humans is expected to open a new avenue of disinfection systems. Details of the inactivation mechanisms of microbes by  $\text{ClO}_2$  gas are also reviewed in this paper.

**CHEMICAL CHARACTERISTICS AND STRUCTURE OF  $\text{ClO}_2$** 

$\text{ClO}_2$  is reddish-yellow gas with an unpleasant odor similar to that of chlorine at room temperature.<sup>[1]</sup>  $\text{ClO}_2$  condenses to a reddish-brown liquid on cooling below  $11^\circ\text{C}$  and freezes at  $-59^\circ\text{C}$ , producing red-orange crystals.<sup>[9]</sup>  $\text{ClO}_2$  is readily soluble in water; 3 grams can be dissolved in 1 liter of water at  $25^\circ\text{C}$ .<sup>[9]</sup>  $\text{ClO}_2$  has one unpaired electron in its molecular orbital and hence is a free radical. Other chemical details of  $\text{ClO}_2$  are presented in Table 1.<sup>[1,9]</sup> Special care is needed to handle pure  $\text{ClO}_2$  gas at high concentrations due to its potential explosiveness and toxicity.<sup>[1,9]</sup> An explosion may result when high concentrations of the gas are exposed to strong light.

**Table 1: Physicochemical characteristics of ClO<sub>2</sub>.**

Characteristic	Value
Molecular weight	67.45
Melting point	-59°C
Boiling point	11°C
Density of liquid	1.642 g/cm <sup>3</sup>
Solubility in water at 25°C	3.01 g/L
Oxidation state of chlorine	+4
Dissociation energy of first Cl-O bond	273 kJ/mol
Dissociation energy of second Cl-O bond	270 kJ/mol
Standard enthalpy of formation (ΔH <sub>f</sub> )	102.6 kJ/mol

The structure of ClO<sub>2</sub> in liquid phase was observed by X-ray diffraction analysis,<sup>[2]</sup> and the gas phase structure was observed by infrared spectroscopy (Table 2).<sup>[10]</sup> Of note, the ClO<sub>2</sub> molecule is bent (C<sub>2v</sub> symmetry) with O-Cl-O angle of 116.1°.<sup>[2]</sup> The Cl-O bond length was 1.46 Å.<sup>[2]</sup> Nielsen and Woltz reported the infrared spectrum of ClO<sub>2</sub> gas, revealing peaks at 290, 445 (ν<sub>2</sub>), 943.2 (ν<sub>1</sub>), 1110, 1888 (2ν<sub>1</sub>), 2040, 2215, 2473, 2967, and 3325 cm<sup>-1</sup>.<sup>[10]</sup> In the solid phase, ClO<sub>2</sub> molecules dimerize, losing their paramagnetic behavior. The dimer (ClO<sub>2</sub>)<sub>2</sub> becomes diamagnetic.<sup>[9]</sup> Shimakura et al. found that in the liquid phase, ClO<sub>2</sub> molecules do not exhibit random orientations, but present a characteristic intermolecular orientation.<sup>[2]</sup>

**Table 2: Chemical structure of ClO<sub>2</sub>.**

Parameter	Value
Cl-O bond length in liquid phase	1.46 Å
Cl-O bond length in gas phase	1.491±0.014 Å
O-Cl-O bond angle in liquid phase	116.1°
O-Cl-O bond angle in gas phase	116.5 ± 2.5°

### GENERATION OF ClO<sub>2</sub> GAS

Numerous methods are available to generate ClO<sub>2</sub> gas. Given that the gas is potentially explosive at high concentrations, it is generally not transported but is generated onsite for use. The most frequently used method of ClO<sub>2</sub> generation involves mixing sodium chlorite (NaClO<sub>2</sub>) with acids or oxidizing agents.<sup>[3]</sup> Sodium chlorite is typically employed as an aqueous solution, and acid is mixed with the solution. ClO<sub>2</sub> generated in the solution is bubbled by air to release it from the solution. For example, the chemical reaction involved in the use of HCl as the acid is 5NaClO<sub>2</sub> + 4HCl → 4ClO<sub>2</sub> + 2H<sub>2</sub>O + 5NaCl. When ClO<sub>2</sub> is generated from chlorine gas (Cl<sub>2</sub>) as a starting material, the chemical reaction is 2NaClO<sub>2</sub> + Cl<sub>2</sub> → 2ClO<sub>2</sub> + 2NaCl.<sup>[3]</sup> ClO<sub>2</sub> is also generated by mixing sodium chlorite with hypochlorous acid (HOCl) following the reaction of 2NaClO<sub>2</sub> + HOCl → NaCl + NaOH + 2ClO<sub>2</sub>. In this reaction, hypochlorous acid is generated by mixing chlorine with water in the reaction of Cl<sub>2</sub> + H<sub>2</sub>O → HOCl + HCl.<sup>[3]</sup>

Electrochemical systems are also employed to generate ClO<sub>2</sub> in situ for use.<sup>[11-13]</sup> In this method, aqueous solution of sodium chlorite is placed in an electrolytic

cell, where ClO<sub>2</sub><sup>-</sup> becomes ClO<sub>2</sub> in an anode and H<sub>2</sub>O becomes hydrogen and OH<sup>-</sup> in a cathode.<sup>[3]</sup> The overall reaction is 2NaClO<sub>2</sub> + 2H<sub>2</sub>O → 2ClO<sub>2</sub> + NaOH + H<sub>2</sub>. ClO<sub>2</sub> in the aqueous solution is stripped from the solution by introducing air into the solution.<sup>[3]</sup> Bai et al. reported a sophisticated method to release ClO<sub>2</sub> gas in a controlled manner.<sup>[14]</sup> ClO<sub>2</sub> gas was generated by adhering two films together. One film is acrylate-based film loaded with sodium chlorite, and the other film is polyvinyl alcohol polymer loaded with tartaric acid. The rate of ClO<sub>2</sub> gas release can be controlled by tailoring film composition and its thickness. The rate of ClO<sub>2</sub> release is accelerated by moisture.<sup>[14]</sup> The researchers noted the usefulness of their system for food packaging. ClO<sub>2</sub> is also generated by exposure of a solution of sodium chlorite to ultraviolet light.<sup>[15,16]</sup> ClO<sub>2</sub> is generated in acidic conditions (pH 3.0-5.0), whereas hypochlorite is generated at alkaline conditions (pH 8.9-10.7).<sup>[15]</sup> Quantum yield of this photochemical reaction irradiated by 253.7 nm ultraviolet light is 0.43 to 0.94 and is maximal at pH 6.<sup>[16]</sup>

### CHEMICAL REACTIONS AND FATE OF ClO<sub>2</sub>

Whether it is an aqueous solution or gas, ClO<sub>2</sub> can react with numerous organic compounds. It is known to react with some free amino acids and amino acid residues in proteins.<sup>[5,17]</sup> For instance, tryptophan and tyrosine, as residues in bovine serum albumin and glucose-6-phosphate dehydrogenase of baker's yeast *Saccharomyces cerevisiae*, were oxidatively modified by an aqueous solution of ClO<sub>2</sub>.<sup>[5]</sup> Furthermore, tryptophan becomes *N*-formylkynurenine, and tyrosine forms 3,4-dihydroxyphenylalanine (DOPA) and 2,4,5-trihydroxyphenylalanine (TOPA).<sup>[5]</sup> Oxygen atoms of ClO<sub>2</sub> were incorporated in the products.<sup>[5]</sup>

Napolitano et al. found that tyrosine, *N*-acetyltyrosine and DOPA react with an aqueous solution of ClO<sub>2</sub>, consuming two molecules of ClO<sub>2</sub> for each reaction.<sup>[17]</sup> In the reaction of tyrosine and *N*-acetyltyrosine, phenoxyl radicals are first generated. Next, a short-lived adduct with a C-OCIO bond at the 3 position of the aromatic ring is generated, ultimately forming dopaquinone and *N*-acetyldopaquinone.<sup>[17]</sup>

The mechanism of the oxidation of tryptophan is proposed as follows.<sup>[18]</sup> Two molecules of ClO<sub>2</sub> react with each molecule of tryptophan. The first molecule

forms a tryptophan radical,  $\text{ClO}_2^-$  and  $\text{H}^+$ . The second molecule reacts with the tryptophan radical and forms a tryptophan- $\text{ClO}$  adduct. Finally the adduct becomes stable *N*-formylkynurenine.<sup>[18]</sup> The oxygen atoms of  $\text{ClO}_2$  are incorporated in the product in this reaction. The amino acid cysteine also reacts with  $\text{ClO}_2$ .<sup>[19]</sup> It is proposed that the reaction involves electron transfer from cysteine anion to  $\text{ClO}_2$  with a subsequent reaction of cysteine radical and  $\text{ClO}_2$  to form a cysteinyl- $\text{ClO}_2$  adduct. The adduct finally forms pH-dependent products: cysteic acid at low pH and cystine at high pH. The tripeptide glutathione (Glu-Cys-Gly) also reacts with  $\text{ClO}_2$ .<sup>[19]</sup>  $\text{ClO}_2$  decomposes in basic aqueous solution via three different pathways.<sup>[20]</sup> One pathway forms  $\text{ClO}_2^-$  and  $\text{O}_2$ . The other two pathways form  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$ . All pathways exhibit a first-order dependence of the reaction with regard to  $\text{OH}^-$ . All the reactions are proposed to proceed by base-assisted electron-transfer mechanisms.<sup>[20]</sup>

Residues after the treatment of objects with  $\text{ClO}_2$  are a particular concern. Basically,  $\text{ClO}_2$  gas is rapidly broken down to chlorate ( $\text{ClO}_3^-$ ) and chlorite ( $\text{ClO}_2^-$ ) ions, which are further converted to chloride ( $\text{Cl}^-$ ) ion.<sup>[21-26]</sup> Kaur et al demonstrated that  $\text{Cl}^-$  and  $\text{ClO}_3^-$  were formed after the treatment of cantaloupes with  $^{36}\text{Cl}$ -labeled  $\text{ClO}_2$  gas.<sup>[22]</sup> They treated cantaloupes with  $5.1 \pm 0.7$  mg/L (1850  $\pm$  254 ppm)  $\text{ClO}_2$  gas for 10 min for fumigation. Then, they measured residues from the rind and flesh of this fruit. They detected  $19.3 \pm 8.0$   $\mu\text{g}$  of  $\text{Cl}^-$  and  $4.8 \pm 2.3$   $\mu\text{g}$  of  $\text{ClO}_3^-$  per gram of rind. They detected  $8.1 \pm 1.0$   $\mu\text{g}$  of  $\text{Cl}^-$  and no  $\text{ClO}_3^-$  per gram of flesh. Given that  $\text{Cl}^-$  is non-toxic, they concluded that fumigation of edible flesh would not pose a health concern.<sup>[22]</sup> Trinetta et al. treated vegetables and fruits with 0.5 mg/L (180 ppm)  $\text{ClO}_2$  gas with 90 to 95% relative humidity for 10 min to disinfected pathological bacteria (*Escherichia coli*, *Listeria monocytogenes* and *Salmonella enterica*). They next rinsed the food surfaces immediately with water to remove any remaining  $\text{ClO}_2$  and byproducts and analyzed the after-rinse water. At 24 h post treatment with  $\text{ClO}_2$  gas, no differences in  $\text{ClO}_2$  residues were noted between control (no  $\text{ClO}_2$  gas treatment) and treated foods such as tomatoes and navel oranges. However,  $\text{ClO}_3^-$  was found in apples. In addition,  $\text{Cl}^-$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  above control values were noted in lettuce.<sup>[23]</sup> Thus, these anions may remain on the surfaces of some agricultural foods treated with  $\text{ClO}_2$  gas if the treatment concentration of the gas is high.

#### MECHANISM OF KINETICS OF ANTIMICROBIAL ACTIVITY OF $\text{ClO}_2$

$\text{ClO}_2$  has strong oxidizing activity presumably due to its free radical properties.<sup>[9]</sup> For instance,  $\text{ClO}_2$  oxidizes tryptophan and tyrosine residues of protein and denatures proteins.<sup>[5,17,27]</sup> In this reaction, oxygen atoms of  $\text{ClO}_2$  are incorporated in the above amino acid residues of proteins, and proteins are denatured.<sup>[5]</sup> The denaturation of proteins and inactivation of enzymes are demonstrated.<sup>[5,28,29]</sup> Finnegan et al. reported degradation

(fragmentation) of bovine serum albumin and aldolase by  $\text{ClO}_2$ .<sup>[28]</sup>

Benarde et al. demonstrated that  $\text{ClO}_2$  kills bacteria by blocking the biosynthesis of bacterial proteins.<sup>[30]</sup> Cho et al. found that  $\text{ClO}_2$  oxidizes bacterial membrane lipids and consequently increases the permeability of the membrane.<sup>[31]</sup> On the other hand, Berg et al. reported that  $\text{ClO}_2$  causes a loss of control of the permeability of  $\text{K}^+$  ion and oxidative damage of the bacterial outer membrane. They concluded that *E. coli* is inactivated by these effects.<sup>[32]</sup> Roller et al. found that dehydrogenase enzymes of *E. coli* are completely inhibited by  $\text{ClO}_2$ , but this effect does not exclusively explain the inactivation mechanism of the bacteria. They suggested that inhibition of protein synthesis might have a contributory lethal effect on the bacteria.<sup>[29]</sup>

Many microorganisms are inactivated by  $\text{ClO}_2$ . Bacteria,<sup>[33-54]</sup> fungi,<sup>[55-58]</sup> viruses<sup>[59-68]</sup> and protozoa<sup>[56-58]</sup> are inactivated.  $\text{ClO}_2$  dissolved in water has long been used to disinfect tap water.<sup>[3]</sup> The antimicrobial activities of  $\text{ClO}_2$  are elicited as a gas. For instance, *Bacillus subtilis*, *S. enterica*, *B. anthracis*, *Francisella tularensis*, *Yersinia pestis*, *E. coli* O157:H7, and *Staphylococcus aureus* are inactivated by  $\text{ClO}_2$  gas.<sup>[4,7,36,37,39,44-46,50,54,69]</sup> Bhagat et al. demonstrated that *S. enterica* inoculated on navel orange surfaces were inactivated to a level of 3.5  $\log_{10}$  reduction of viability with 0.1 mg/L (36 ppm)  $\text{ClO}_2$  gas for 12 min at 22°C with 90-95 % relative humidity.<sup>[50]</sup> A mixture of *S. enterica*, *E. coli* and *L. monocytogenes* spot-inoculated on the surface of tomatoes, cantaloupes and strawberries were treated with 10 mg/L  $\text{ClO}_2$  (3600 ppm) gas for 180 s, and a 3-5  $\log_{10}$  reduction of viability was reported.<sup>[35]</sup>

The inactivation activity of the  $\text{ClO}_2$  gas is elicited against the bacteria not only in their floating state in the air, but also in their attachment state on solid objects.<sup>[69]</sup> Li et al. found that spore-forming bacteria, *B. subtilis* var. *niger* attached to pieces of metal, plastic, and glass were inactivated by 800 ppm (2.2 mg/L)  $\text{ClO}_2$  gas for 3 h to levels of 1.8 to 6.64  $\log_{10}$  reduction.<sup>[69]</sup> The results clearly indicated sporocidal activity of the gas. Interestingly, the inactivation activity is dependent on pre-humidification treatment of the test pieces.<sup>[69]</sup> Viruses are also inactivated by the gas. For instance, influenza virus, feline calicivirus, human herpesvirus, and canine distemper virus are inactivated by  $\text{ClO}_2$  gas.<sup>[6,27,64,70,71]</sup>

Wang et al. extensively studied the kinetics of the inactivation of *B. subtilis* spores and *S. albus* inoculated on a piece of filter paper by  $\text{ClO}_2$  gas.<sup>[46]</sup> They fitted the rate of the kill of the bacteria using a first-order kinetic model using a function of  $\log_{10}(N/N_0) = -kt$ , where  $N_0$  is the initial number of cells,  $N$  is the number of surviving bacteria after time  $t$  (min) of  $\text{ClO}_2$  gas exposure, and  $k$  ( $\text{min}^{-1}$ ) is the rate constant. In the case of *B. subtilis* spores, the rate constant  $k$  is  $0.09 \pm 0.01$   $\text{min}^{-1}$  ( $n = 6$ ) at

1 mg/L gas concentration with 70% relative humidity at 22 to 24°C, and this value increases to  $0.21 \pm 0.02 \text{ min}^{-1}$  at 5 mg/L.<sup>[46]</sup> In the case of *S. albus*,  $k$  was  $0.15 \pm 0.01 \text{ min}^{-1}$  at 2 mg/L gas with 70% relative humidity and  $0.32 \pm 0.02 \text{ mg/L}$  at 5 mg/L gas.<sup>[46]</sup> Interestingly, the rate constant  $k$  decreases to  $0.04 \pm 0.01 \text{ min}^{-1}$  at 2 mg/L gas with 30% relative humidity, whereas it increases to  $0.66 \pm 0.04 \text{ min}^{-1}$  with 90% relative humidity. Thus, the rate of killing increases along with the increase in relative humidity. The same trend is also noted in the case of *B. subtilis* spores.<sup>[46]</sup> The augmentation of the inactivation activity of ClO<sub>2</sub> gas upon the increase in relative humidity was also reported regarding *S. enteritidis* inoculated on eggshells.<sup>[45]</sup> Of note, the same trend of the effect of humidity was also reported regarding the inactivation of feline calicivirus. Morino et al. found that feline calicivirus placed on a glass surface and treated with 0.26 ppm (0.72 µg/L) ClO<sub>2</sub> gas for 24 h at 20°C was inactivated to 1.0 log<sub>10</sub> reduction ( $n = 4$ ) with 45 to 55% relative humidity, whereas it was inactivated to 6.3 log<sub>10</sub> reduction with 75 to 85% relative humidity.<sup>[70]</sup>

### CLO<sub>2</sub> GAS AS A FUMIGANT

Previously high-concentration ClO<sub>2</sub> gas was frequently used as a fumigant to inactivate various microbes. For instance, Park and Kang demonstrated that *E. coli*, *S. typhimurium* and *L. monocytogenes* inoculated on spinach leaves and tomato surfaces were inactivated by 5 or 10 ppm (28 µg/L) ClO<sub>2</sub> gas.<sup>[33]</sup> *S. enterica*, *E. coli* O157:H7 and *L. monocytogenes* spotted on the surface of crops (tomatoes, cantaloupes and strawberries) were treated with 10 mg/L (3600 ppm) gas of ClO<sub>2</sub> for 180 s.<sup>[35]</sup> In this experiment, a 5-log<sub>10</sub> reduction in colony forming unit (CFU) was noted in *S. enterica* in all crops. In contrast, a 3-log<sub>10</sub> reduction in CFU was noted in *E. coli* and *L. monocytogenes*, indicating that the latter two are more resistant to ClO<sub>2</sub> gas.<sup>[35]</sup> A 3-log<sub>10</sub> CFU reduction of *S. enterica* was also reported for mung bean sprouts at 0.5 mg/L (180 ppm) ClO<sub>2</sub> gas for 15 min.<sup>[37]</sup> Of note, the inactivation activity of ClO<sub>2</sub> gas was also found on bacterial spores.<sup>[34,36,38,42,46,47,72,75]</sup> Jenk and Woodworth reported that spore-forming bacteria *B. subtilis* planted in the artificial organs were reproducibly sterilized with 30-min dwell time with 30 mg/L ClO<sub>2</sub> gas (10900 ppm) with 80 to 85% relative humidity at 30°C.<sup>[72]</sup> The D value (time required for 90% spore inactivation) was 4.4 min.<sup>[72]</sup> Lowe et al. found that 362 – 695 ppm ClO<sub>2</sub> gas maintained at exposures of 756 ppm-hours with 65% relative humidity achieved inactivation of *B. anthracis* and *Mycobacterium smegmatis*. The reduction of viability was greater than 6 log<sub>10</sub>.<sup>[34]</sup>

### TOXICITY STUDY OF CLO<sub>2</sub> GAS

High-concentration ClO<sub>2</sub> gas is toxic against numerous animals including human. Paulet and Desbrousses performed a toxicological study. Rats exposed to 10 ppm (28 µg/L) ClO<sub>2</sub> gas for 2 h/day for the 30-day period exhibited nasal discharge, red eyes, localized bronchopneumonia with desquamation of the alveolar

epithelium and an increase in leukocytes.<sup>[74]</sup> The same group also performed other experiment using rats. They exposed rats to 1 ppm (2.8 µg/L) ClO<sub>2</sub> gas for 5 h/day for 5 days/week during a 2-month period. They found vascular congestion and peribronchiolar edema in the lungs of the rats.<sup>[75]</sup> However, Ogata et al.<sup>[8]</sup> could not reproduce their pathological findings in rats exposed to the experimental conditions exactly as described in the paper of Paulet and Desbrousses.<sup>[75]</sup> They concluded that the most likely reason for this discrepancy might involve the fine controls of gas concentrations.<sup>[8]</sup> As noted by Ogata et al.,<sup>[8]</sup> fine controls of ClO<sub>2</sub> gas concentrations at such low levels might have been quite difficult to achieve at the time of Paulet and Desbrousses<sup>[75]</sup> as a gas generator with a sophisticated control system was not available.<sup>[8]</sup>

Akamatsu et al. demonstrated the rats exposed to 0.1 ppm ClO<sub>2</sub> gas for 24 h/day and 7 days/week for a period of 6 months were completely healthy at the end of the experiment.<sup>[76]</sup> Dalhamn conducted ClO<sub>2</sub> gas inhalation study on rats. Exposure to 260 ppm (720 µg/L) ClO<sub>2</sub> gas for 2 h resulted in ocular discharge, epistaxis, pulmonary edema, circulatory engorgement and death.<sup>[77]</sup> In contrast, exposure to 0.1 ppm (0.28 µg/L) ClO<sub>2</sub> gas for 5 h/day during a 10-week period did not cause any pathological effect, and he concluded that this level is NOAEL (no-observed-adverse-effect level).<sup>[77]</sup> Exposure of rats to 10 ppm (28 µg/L) ClO<sub>2</sub> gas for 4 h/day during a 2-week period caused respiratory tract irritation, and he concluded that this level is LOAEL (lowest-observed-adverse-effect level).<sup>[77]</sup>

### GOVERNMENTAL REGULATIONS OF CLO<sub>2</sub> GAS CONCENTRATIONS

Given that high-concentration ClO<sub>2</sub> gas and liquid are explosive and toxic to animals as mentioned above, several governmental regulations have been implemented in some countries. American OSHA (Occupational Safety and Health Administration) states that the 8-hour time-weighted average of permissible exposure level of the ClO<sub>2</sub> gas is 0.1 ppm (0.28 µg/L).<sup>[78]</sup> The American Conference of Governmental Industry Hygienist (ACGIH) also states 0.1 ppm as a permissible level for workers working 40 hours per week and 8 hours a day.<sup>[79]</sup> NIOSH (National Institute for Occupational Safety and Health) of USA states that the permissible average 10-hour exposure level is 0.1 ppm to humans.<sup>[80]</sup> Taken together, exposure to less than 0.1 ppm ClO<sub>2</sub> gas appears to be safe for humans. Thus, it would be concluded that long-term exposure to ClO<sub>2</sub> gas at or below 0.1 ppm would be allowable to humans.

### ANTIMICROBIAL ACTIVITY OF CLO<sub>2</sub> GAS AT EXTREMELY LOW CONCENTRAIONS

Ogata and Shibata first reported the effect of extremely low-concentration ClO<sub>2</sub> gas at a level of 0.03 ppm (0.084 µg/L) against influenza virus in an animal experiment<sup>[6]</sup> using a sophisticated machine to generate and deliver ClO<sub>2</sub> gas at finely controlled concentrations.<sup>[7]</sup> The gas



concentration was precisely controlled and accurately monitored during the study as demonstrated by recently published paper.<sup>[81]</sup> They found that the lethal activity of influenza A virus aerosol exposed to mice was dramatically reduced when 0.03 ppm ClO<sub>2</sub> gas was present simultaneously with the virus aerosol. All the virus-challenged mice were alive and appeared quite healthy during and after the exposure of the virus when ClO<sub>2</sub> gas was concomitantly present.<sup>[6]</sup> This result suggests the potential usefulness of the gas to protect human diseases caused by floating microbes in a room. A crucial point of this result is that evacuation of people from the room would not be required during the exposure to the gas because the concentration of the ClO<sub>2</sub> gas employed is extremely low, i.e., below the permissible exposure concentration to human as mentioned above.<sup>[79,80,82]</sup> Thus, the exposure is *not fumigation*. Currently there is no useful and reliable measure to protect humans from infection by floating microbes without requiring evacuation in closed or semi-closed spaces, such as an airplane cabin or a spacecraft. The prevention of airborne microbe infection by the extremely low-concentration of ClO<sub>2</sub> gas will open new avenues in the field of public health, e.g., prevention of highly pathogenic and transmissible H5N1 influenza virus.<sup>[83]</sup> The use of 0.03 ppm ClO<sub>2</sub> gas is also useful in prevention of mosquito-related infective diseases, such as malaria and dengue fever, given that this concentration of ClO<sub>2</sub> gas has a repellent effect against mosquitoes.<sup>[84]</sup>

### CONCLUSION

Exposure to extremely low concentrations of ClO<sub>2</sub> gas, i.e., at or below 0.1 ppm (0.28 µg/L), has no harmful effect on animals, whereas 0.03 to 0.1 ppm still has inactivation activities against bacteria and virus. Such concentrations of ClO<sub>2</sub> gas could be used without requiring evacuation of people to prevent infections by microbes floating in air in closed or semi-closed spaces, such as in the cabins of aircrafts, living rooms and spacecraft. This effect of ClO<sub>2</sub> gas can be used to prevent the spread of infectious diseases, such as highly pathogenic H5N1 influenza virus, by increasing the quality of indoor air. Currently, such a disinfectant is not commercially available. To the best of our knowledge, the extremely low concentrations of ClO<sub>2</sub> are the only measure to prevent the infection by airborne microbes in the presence of humans.

### ACKNOWLEDGEMENTS

This work was supported by Taiko Pharmaceutical Co., Ltd.

### Declaration of Interest

The author is an employee of Taiko Pharmaceutical Co., Ltd.

### REFERENCES

- O'Neil MJ. The Merck Index. 14th ed., Whitehouse Station; Merck, 2006.

- Shimakura H, Ogata N, Kawakita Y, Ohara K, Takeda S. Determination of the structure of liquids containing free radical molecules: Inter-molecular correlations in liquid chlorine dioxide. *Mol Phys*, 2013; 111(8): 1015-22.
- Gates DJ. The Chlorine Dioxide Handbook Water Disinfection Series. Denver; American Water Works Association, 1998.
- Morino H, Fukuda T, Miura T, Shibata T. Effect of low-concentration chlorine dioxide gas against bacteria and viruses on a glass surface in wet environments. *Lett Appl Microbiol*, 2011; 53(6): 628-34.
- Ogata N. Denaturation of protein by chlorine dioxide: oxidative modification of tryptophan and tyrosine residues. *Biochemistry*, 2007; 46(16): 4898-911.
- Ogata N, Shibata T. Protective effect of low-concentration chlorine dioxide gas against influenza A virus infection. *J Gen Virol*, 2008; 89(Pt 1): 60-7.
- Ogata N, Sakasegawa M, Miura T, Shibata T, Takigawa Y, Taura K, Taguchi K, Matsubara K, Nakahara K, Kato D, Sogawa K, Oka H. Inactivation of airborne bacteria and viruses using extremely low concentrations of chlorine dioxide gas. *Pharmacology*, 2016; 97(5-6): 301-6.
- Ogata N, Koizumu T, Ozawa F. Ten-week whole-body inhalation toxicity study of chlorine dioxide gas in rats. *J Drug Metab Toxicol*, 2013; 4(2): 143.
- Wiberg N. Holleman-Wiberg's Inorganic Chemistry. San Diego; Academic Press, 2001.
- Nielsen AH, Woltz PJH. The infrared spectrum of chlorine dioxide. *J Chem Phys.*, 1952; 20(12): 1878-83.
- Cowley G. ClO<sub>2</sub> in the pulp industry: A novel electrochemical cell for generation of ClO<sub>2</sub> from aqueous sodium chlorite using the Perstraction™ process. In: Proc EMA, USEPA, and AWWARF 3rd International Symposium on ClO<sub>2</sub>. New Orleans, 1995.
- Kaczur JJ, Cawfield DW. Chlorous acid, chlorites and chlorine dioxide (ClO<sub>2</sub>, HClO<sub>2</sub>). In: Othmer D (ed): Kirk-Othmer Encyclopedia of Chemical Technology. 4th ed., New York; Wiley Interscience, 1993; 5: 968-97.
- Griese MH, Kaczur JJ, Gordon G. Combining methods for the reduction of oxychlorine residuals in drinking water. *J Am Water Works Assoc*, 1992; 84(11): 69-77.
- Bai Z, Cristancho DE, Rachford AA, Reder AL, Williamson A, Grzesiak AL. Controlled release of antimicrobial ClO<sub>2</sub> gas from a two-layer polymeric film system. *J Agric Food Chem*, 2016; 64(45): 8647-52.
- Kujrai C, Fujita I. Photochemistry of sodium chlorite in solution, with reference to chlorite bleaching. *J Soc Dyers Colourists*, 1962; 78(2): 80-9.
- Cosson H, Ernst WR. Photodecomposition of chlorine dioxide and sodium chlorite in aqueous

- solution by irradiation with ultraviolet light. *Ind Eng Chem Res.*, 1994; 33(6): 1468-75.
17. Napolitano MJ, Green BJ, Nicoson JS, Margerum DW. Chlorine dioxide oxidations of tyrosine, *N*-acetyltyrosine, and dopa. *Chem Res Toxicol*, 2005; 18(3): 501-8.
  18. Stewart DJ, Napolitano MJ, Bakhmutova-Albert EV, Margerum DW. Kinetics and mechanisms of chlorine dioxide oxidation of tryptophan. *Inorg Chem*, 2008; 47(5): 1639-47.
  19. Ison A, Odeh IN, Margerum DW. Kinetics and mechanisms of chlorine dioxide and chlorite oxidations of cysteine and glutathione. *Inorg Chem.*, 2006; 45(21): 8768-75.
  20. Odeh IN, Francisco JS, Margerum DW. New pathways for chlorine dioxide decomposition in basic solution. *Inorg Chem.*, 2002; 41(24): 6500-6.
  21. Gómez-López VM, Rjkovic A, Ragaert P, Smigic N, Devileghere F. Chlorine dioxide for minimally processed produce preservation: a review. *Trends Food Sci Tech.*, 2009; 20(1): 17-26.
  22. Kaur S, Smith DJ, Morgan MT. Chloroxyanion residue quantification in cantaloupes treated with chlorine dioxide gas. *J Food Prot.*, 2015; 78(9): 1708-18.
  23. Trinetta V, Vaidya N, Linton R, Morgan M. Evaluation of chlorine dioxide gas residues on selected food produce. *J Food Sci.*, 2011; 76(1): T11-5.
  24. Smith DJ, Ernst W, Giddings JM. Distribution and chemical fate of <sup>36</sup>Cl-chlorine dioxide gas during the fumigation of tomatoes and cantaloupe. *J Agric Food Chem.*, 2014; 62(48): 11756-66.
  25. Smith DJ, Ernst W, Herges GR. Chloroxyanion residues in cantaloupe and tomatoes after chlorine dioxide gas sanitation. *J Agric Food Chem.*, 2015; 63(43): 9640-9.
  26. Smith DJ, Giddings JM, Herges GR, Ernst W. Distribution, identification, and quantification of residues after treatment of ready-to-eat salami with <sup>36</sup>Cl-labeled or nonlabeled chlorine dioxide gas. *J Agric Food Chem.*, 2016; 64(44): 8454-62.
  27. Ogata N. Inactivation of influenza virus haemagglutinin by chlorine dioxide: oxidation of the conserved tryptophan 153 residue in the receptor-binding site. *J Gen Virol*, 2012; 93: 2558-63.
  28. Finnegan M, Linley E, Denyer SP, McDonnell G, Simons C, Maillard JY. Mode of action of hydrogen peroxide and other oxidizing agents: differences between liquid and gas forms. *J Antimicrob Chemother*, 2010; 65(10): 2108-15.
  29. Roller SD, Olivieri VP, Kawata K. Mode of bacterial inactivation by chlorine dioxide. *Water Res.*, 1980; 14(6): 635-41.
  30. Benarde MA, Snow WB, Olivieri VP, Davidson B. Kinetics and mechanism of bacterial disinfection by chlorine dioxide. *Appl Microbiol*, 1967; 15(2): 257-65.
  31. Cho M, Kim J, Kim JY, Yoon J, Kim JH. Mechanisms of *Escherichia coli* inactivation by several disinfectants. *Water Res.*, 2010; 44(11): 3410-8.
  32. Berg JD, Roberts PV, Matin A. Effect of chlorine dioxide on selected membrane functions of *Escherichia coli*. *J Appl Bacteriol*, 1986; 60(3): 213-20.
  33. Park SH, Kang DH. Combination treatment of chlorine dioxide gas and aerosolized sanitizer for inactivating foodborne pathogens on spinach leaves and tomatoes. *Int J Food Microbiol*, 2015; 207: 103-8.
  34. Lowe JJ, Hewlett AL, Iwen PC, Smith PW, Gibbs SG. Evaluation of ambulance decontamination using gaseous chlorine dioxide. *Prehosp Emerg Care*, 2013; 17(3): 401-8.
  35. Trinetta V, Linton RH, Morgan MT. The application of high-concentration short-time chlorine dioxide treatment for selected specialty crops including Roma tomatoes (*Lycopersicon esculentum*), cantaloupes (*Cucumis melo* ssp. *melo* var. *cantaloupensis*) and strawberries (*Fragaria × ananassa*). *Food Microbiol*, 2013; 34(2): 296-302.
  36. Lowe JJ, Gibbs SG, Iwen PC, Smith PW, Hewlett AL. Decontamination of a hospital room using gaseous chlorine dioxide: *Bacillus anthracis*, *Francisella tularensis*, and *Yersinia pestis*. *J Occup Environ Hyg*, 2013; 10(10): 533-9.
  37. Prodduk V, Annous BA, Liu L, Yam KL. Evaluation of chlorine dioxide gas treatment to inactivate *Salmonella enterica* on mungbean sprouts. *J Food Prot*, 2014; 77(11): 1876-81.
  38. Rastogi VK, Ryan SP, Wallace L, Smith LS, Shah SS, Martin GB. Systematic evaluation of the efficacy of chlorine dioxide in decontamination of building interior surfaces contaminated with anthrax spores. *Appl Environ Microbiol*, 2010; 76(10): 3343-51.
  39. Bhagat A, Mahmoud BS, Linton RH. Inactivation of *Salmonella enterica* and *Listeria monocytogenes* inoculated on hydroponic tomatoes using chlorine dioxide gas. *Foodborne Pathog Dis.*, 2010; 7(6): 677-85.
  40. Vaid R, Linton RH, Morgan MT. Comparison of inactivation of *Listeria monocytogenes* within a biofilm matrix using chlorine dioxide gas, aqueous chlorine dioxide and sodium hypochlorite treatments. *Food Microbiol*, 2010; 27(8): 979-84.
  41. Kuroyama I, Osato S, Nakajima S, Kubota R, Ogawa T. Environmental monitoring and bactericidal efficacy of chlorine dioxide gas in a dental office. *Biocontrol Sci.*, 2010; 15(3): 103-9.
  42. Shirasaki Y, Matsuura A, Uekusa M, Ito Y, Hayashi T. A study of the properties of chlorine dioxide gas as a fumigant. *Exp Anim*, 2016; 65(3): 303-10.
  43. Page N, González-Buesa J, Ryser ET, Harte J, Almenar E. Interactions between sanitizers and packaging gas compositions and their effects on the safety and quality of fresh-cut onions (*Allium cepa* L.). *Int J Food Microbiol*, 2016; 218: 105-13.

44. Kim HG, Song KB. Combined treatment with low concentrations of aqueous and gaseous chlorine dioxide inactivates *Escherichia coli* O157:H7 and *Salmonella typhimurium* inoculated on paprika. *J Microbiol Biotechnol*, 2016; doi:10.4014/jmb.1611.11038.
45. Kim H, Yum B, Yoon SS, Song KJ, Kim JR, Myeong D, Chang B, Choe NH. Inactivation of *Salmonella* on eggshells by chlorine dioxide gas. *Korean J Food Sci Anim Resour*, 2016; 36(1): 100-8.
46. Wang T, Wu J, Qi J, Hao L, Yi Y, Zhang Z. Kinetics of inactivation of *Bacillus subtilis* subsp. *niger* spores and *Staphylococcus albus* on paper by chlorine dioxide gas in an enclosed space. *Appl Environ Microbiol*, 2016; 82(10): 3061-9.
47. Wang T, Qi J, Wu J, Hao L, Yi Y, Lin S, Zhang Z. Response surface modeling for the inactivation of *Bacillus subtilis* subsp. *niger* spores by chlorine dioxide gas in an enclosed space. *J Air Waste Manag Assoc*, 2016; 66(5): 508-17.
48. Li YJ, Zhu N, Jia HQ, Wu JH, Yi Y, Qi JC. Decontamination of *Bacillus subtilis* var. *niger* spores on selected surfaces by chlorine dioxide gas. *J Zhejiang Univ Sci B*, 2012; 13(4): 254-60.
49. Trinetta V, Vaidya N, Linton R, Morgan M. A comparative study on the effectiveness of chlorine dioxide gas, ozone gas and e-beam irradiation treatments for inactivation of pathogens inoculated onto tomato, cantaloupe and lettuce seeds. *Int J Food Microbiol*, 2011; 146(2): 203-6.
50. Bhagat A, Mahmoud BS, Linton RH. Effect of chlorine dioxide gas on *Salmonella enterica* inoculated on navel orange surfaces and its impact on the quality attributes of treated oranges. *Foodborne Pathog Dis.*, 2011; 8(1): 77-85.
51. Kaur S, Smith DJ, Morgan MT. Chloroxyanion residue quantification in cantaloupes treated with chlorine dioxide gas. *J Food Prot.*, 2015; 78(9):1708-18.
52. Lowe JJ, Gibbs SG, Iwen PC, Smith PW, Hewlett AL. Impact of chlorine dioxide gas sterilization on nosocomial organism viability in a hospital room. *Int J Environ Res Public Health*, 2013; 10(6): 2596-605.
53. Pottage T, Macken S, Giri K, Walker JT, Bennett AM. Low-temperature decontamination with hydrogen peroxide or chlorine dioxide for space applications. *Appl Environ Microbiol*, 2012; 78(12): 4169-74.
54. Annous BA, Burke A. Development of combined dry heat and chlorine dioxide gas treatment with mechanical mixing for inactivation of *Salmonella enterica* serovar Montevideo on mung bean seeds. *J Food Prot*, 2015; 78(5): 868-72.
55. Sun X, Bai J, Ference C, Wang Z, Zhang Y, Narciso J, Zhou K. Antimicrobial activity of controlled-release chlorine dioxide gas on fresh blueberries. *J Food Prot*, 2014; 77(7): 1127-32.
56. Erickson MC, Ortega YR. Inactivation of protozoan parasites in food, water, and environmental systems. *J Food Prot*, 2006; 69(11): 2786-808.
57. Ortega YR, Mann A, Torres MP, Cama V. Efficacy of gaseous chlorine dioxide as a sanitizer against *Cryptosporidium parvum*, *Cyclospora cayetanensis*, and *Encephalitozoon intestinalis* on produce. *J Food Prot*, 2008; 71(12): 2410-4.
58. Winiiecka-Krusnell J, Linder E. Cysticidal effect of chlorine dioxide on *Giardia intestinalis* cysts. *Acta Trop*, 1998; 70(3): 369-72.
59. Miura T, Shibata T. Antiviral effect of chlorine dioxide against influenza virus and its application for infection control. *Open Antimicrobial Agents J.*, 2010; 2: 71-8.
60. Taylor GR, Butler M. A comparison of the virucidal properties of chlorine, chlorine dioxide, bromine chloride and iodine. *J Hyg (Lond)*, 1982; 89(2): 321-8.
61. Junli H, Li W, Nenqi R, Li LX, Fun SR, Guanle Y. Disinfection effect of chlorine dioxide on viruses, algae and animal planktons in water. *Water Res.*, 1997; 31(3): 455-60.
62. Chen YS, Vaughn JM. Inactivation of human and simian rotaviruses by chlorine dioxide. *Appl Environ Microbiol*, 1990; 56(5): 1363-6.
63. Wigginton KR, Pecson BM, Sigstam T, Bosshard F, Kohn T. Virus inactivation mechanisms: impact of disinfections on virus function and structural integrity. *Environ Sci Technol*, 2012; 46(21): 12069-78.
64. Sanekata T, Fukuda T, Miura T, Morino H, Lee C, Maeda K, Araki K, Otake T, Kawahata T, Shibata T. Evaluation of the antiviral activity of chlorine dioxide and sodium hypochlorite against feline calicivirus, human influenza virus, measles virus, canine distemper virus, human herpesvirus, human adenovirus, canine adenovirus and canine parvovirus. *Biocontrol Sci.*, 2010; 15(2): 45-9.
65. Watamoto T, Egusa H, Sawase T, Yatani H. Clinical evaluation of chlorine dioxide for disinfection of dental instruments. *Int J Prosthodont*, 2013; 26(6): 541-4.
66. Lowe JJ, Hewlett AL, Iwen PC, Smith PW, Gibbs SG. Surrogate testing suggests that chlorine dioxide gas exposure would not inactivate Ebola virus contained in environmental blood contamination. *J Occup Environ Hyg*, 2015; 12(9): D211-5.
67. Yeap JW, Kaur S, Lou F, DiCaprio E, Morgan M, Linton R, Li J. Inactivation kinetics and mechanism of a human norovirus surrogate on stainless steel coupons via chlorine dioxide gas. *Appl Environ Microbiol*, 2016; 82(1): 116-23.
68. Kingsley, DH, Vincent EM, Meade GK, Watson CL, Fan X. Inactivation of human norovirus using chemical sanitizers. *Int J Food Microbiol*, 2014; 171: 94-9.
69. Li Y, Zhu N, Jia HQ, Wu JH, Yi Y, Qi JC. Decontamination of *Bacillus subtilis* var. *niger*

- spores on selected surfaces by chlorine dioxide gas. *J Zhejiang Univ Sci B*, 2012; 13(4): 254-60.
70. Morino H, Fukuda T, Miura T, Lee C, Shibata T, Sanekata T. Inactivation of feline calicivirus, a norovirus surrogate, by chlorine dioxide gas. *Biocontrol Sci.*, 2009; 14(4): 147-53.
  71. Morino H, Koizumi T, Miura T, Fukuda T, Shibata T. Inactivation of feline calicivirus by chlorine dioxide gas-generating gel. *Yakugaku Zasshi*, 2012; 133(9): 1017-22.
  72. Jeng DK, Woodworth AG. Chlorine dioxide gas sterilization of oxygenators in an industrial scale sterilizer: a successful model. *Artificial Organs*, 1990; 14(5): 361-8.
  73. Rastogi VK, Wallace L, Smith LS, Ryan SP, Martin B. Quantitative method to determine sporicidal decontamination of building surfaces by gaseous fumigants, and issues related to laboratory-scale studies. *Appl Environ Microbiol*, 2009; 75(11): 3688-94.
  74. Paulet G, Desbrousses S. Effect of a weak concentration of chlorine dioxide on laboratory animals. *Arch Mal Prof*, 1970; 31(3): 97-106.
  75. Paulet G, Desbrousses S. On the toxicology of chlorine dioxide. *Arch Mal Prof*, 1972; 33(1): 59-61.
  76. Akamatsu A, Lee C, Morino H, Miura T, Ogata N, Shibata T. Six-month low level chlorine dioxide gas inhalation toxicity study with two-week recovery period in rats. *J Occup Med Toxicol*, 2012; 7: 2.
  77. Dalhamn T. Chlorine dioxide: toxicity in animal experiments and industrial risks. *Arch Ind Health*, 1957; 15(2): 101-7.
  78. Occupational Safety and Health Administration (OSHA), United States of Labor. Annotated PELs, <https://www.osha.gov/dsg/annotated/pels/tablez-1.html>.
  79. ACGIH. 1994-1995 Threshold limit values for chemical substances and physical agents and biological exposure indices. American Conference of Governmental Industrial Hygienists, Cincinnati, OH, 1994.
  80. NIOSH. Recommendation for occupational safety and health: Compendium of policy documents and statements. U. S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 92-100, Cincinnati, OH, Jan., 1992.
  81. Ogata N, Sogawa K, Takigawa Y, Shibata T. Generation and measurement of chlorine dioxide gas at extremely low concentrations in a living room: implication for preventing airborne microbial infectious diseases. *Pharmacology*, 2017; 99(3-4): 114-20.
  82. Harfoot R, Webby RJ. H5 influenza, a global update. *J Microbiol*, 2017; 55(3): 196-203.
  83. Matsuoka H, Ogata N. Inhibition of malaria infection and repellent effect against mosquitoes by chlorine dioxide. *Med Entomol Zool*, 2013; 64(4): 203-7.